

Table A2. Cases of HIV infection (not AIDS) for male adults and adolescents, by exposure category and race/ethnicity, reported through December 2002—39 areas with confidential name-based HIV infection reporting

Exposure category	White, not Hispanic				Black, not Hispanic				Hispanic			
	2002		Cumulative through 2002 ^a		2002		Cumulative through 2002 ^a		2002		Cumulative through 2002 ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact	5,986	66	36,920	64	2,930	30	19,135	32	1,917	42	7,171	41
Injection drug use	506	6	4,543	8	1,039	11	9,753	16	589	13	2,838	16
Male-to-male sexual contact and injection drug use	393	4	4,216	7	231	2	2,690	4	105	2	648	4
Hemophilia/coagulation disorder	17	0	349	1	6	0	99	0	4	0	19	0
Heterosexual contact	282	3	1,781	3	1,185	12	7,490	12	336	7	1,298	7
Sex with injection drug user	67	1	438	1	159	2	1,285	2	39	1	215	1
Sex with person with hemophilia	2	0	5	0	0	0	13	0	2	0	3	0
Sex with HIV-infected transfusion recipient	3	0	24	0	11	0	64	0	3	0	8	0
Sex with HIV-infected person, risk not specified	210	2	1,314	2	1,015	10	6,128	10	292	6	1,072	6
Receipt of blood transfusion, blood components, or tissue	19	0	215	0	21	0	206	0	16	0	42	0
Other/risk not reported or identified	1,888	21	9,411	16	4,322	44	20,872	35	1,580	35	5,460	31
Total	9,091	100	57,435	100	9,734	100	60,245	100	4,547	100	17,476	100

Exposure category	Asian/Pacific Islander				American Indian/Alaska Native				Totals			
	2002		Cumulative through 2002 ^a		2002		Cumulative through 2002 ^a		2002		Cumulative through 2002 ^a	
	No.	%	No.	%	No.	%	No.	%	No.	%	No.	%
Male-to-male sexual contact	106	52	416	52	52	58	411	55	10,991	46	64,331	47
Injection drug use	8	4	37	5	7	8	88	12	2,149	9	17,312	13
Male-to-male sexual contact and injection drug use	2	1	15	2	7	8	99	13	738	3	7,693	6
Hemophilia/coagulation disorder	0	0	2	0	0	0	1	0	27	0	473	0
Heterosexual contact:	17	8	52	7	5	6	43	6	1,825	8	10,689	8
Sex with injection drug user	2	1	8	1	1	1	15	2	268	1	1,964	1
Sex with person with hemophilia	0	0	0	0	0	0	0	0	4	0	21	0
Sex with HIV-infected transfusion recipient	0	0	2	0	0	0	0	0	17	0	98	0
Sex with HIV-infected person, risk not specified	15	7	42	5	4	4	28	4	1,536	6	8,606	6
Receipt of blood transfusion, blood components, or tissue	0	0	3	0	0	0	2	0	56	0	473	0
Other/risk not reported or identified	72	35	271	34	18	20	107	14	7,880	33	37,046	27
Total	205	100	796	100	89	100	751	100	23,666	100	138,017	100

Note. Includes only persons with HIV infection that has not progressed to AIDS.

Since 2002, the following 39 areas have had laws or regulations requiring confidential name-based HIV infection reporting: Alabama, Alaska, Arizona, Arkansas, Colorado, Connecticut, Florida, Idaho, Indiana, Iowa, Kansas, Louisiana, Michigan, Minnesota, Mississippi, Missouri, Nebraska, Nevada, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Utah, Virginia, West Virginia, Wisconsin, Wyoming, American Samoa, Guam, Northern Mariana Islands, and the U.S. Virgin Islands.

Connecticut has confidential name-based HIV infection reporting only for pediatric cases.

Florida (since July 1997) has had confidential name-based HIV infection reporting only for new diagnoses.

Pennsylvania (October 2002) implemented confidential name-based HIV infection reporting only in areas outside the city of Philadelphia.

Texas (February 1994 through December 1998) reported only pediatric HIV infection cases.

^a Includes persons with a diagnosis of HIV infection (not AIDS), reported from the beginning of the epidemic through 2002. Cumulative total includes 1314 males of unknown or multiple race.